

Effect of one-ion L - S coupling on magnetic properties of a correlated spin-orbit system

Zu-Jian Ying^{1,2,3}, Xi-Wen Guan⁴, Angela Foerster¹, Itzhak Roditi³, Bin Chen²

1. Instituto de Física da UFRGS, Av. Bento Gonçalves, 9500, Porto Alegre, 91501-970, Brasil

2. Hangzhou Teachers College, Hangzhou 310012, China

3. Centro Brasileiro de Pesquisas Físicas, Rua Dr. Xavier Sigaud 150, 22290-180 Rio de Janeiro, RJ, Brasil

4. Department of Theoretical Physics, Research School of Physical Sciences and Engineering, and Centre for Mathematics and its Applications, Mathematical Sciences Institute, Australian National University, Canberra ACT 0200, Australia

()

By introducing a basis for a novel realization of the $SU(4)$ Lie algebra, we exactly solve a spin-orbital chain with one-ion L - S coupling (OILSC) via the Bethe ansatz (BA) approach. In the context of different Landé g factors of the spin and orbital sectors, the OILSC results in rich and novel quantum phase transitions. Some accurate analytical expressions for the critical fields are obtained. Both spin ordering and orbital ordering are found in a gapped singlet phase. The system exhibits many interesting phenomena such as nonvanishing magnetization of the singlet phase, multi-entrance of the singlet in the ground state when the field varies, unsymmetric magnetization in two-component phase and magnetization crossover for different OILSC.

Strongly-correlated electron systems in the presence of orbital degeneracy have attracted much interest due to experimental advances in transition-metal compounds [1], and the orbital ordering as well as the orbital density wave has been observed in a family of manganites [2]. Among the experimental findings are materials related to spin-orbital systems in one dimension, such as tetrahis(dimethylamino) ethylene- C_{60} [3], artificial quantum dot arrays [4], $Na_2Ti_2Sb_2O$ and NaV_2O_5 degenerate chains [5], and so forth. A well-known spin-orbital model is the $SU(4)$ model [6] which is exactly solvable in one dimension [7,8]. The competition of orbital degree of freedom with the spin results in three quantum phase transitions (QPT's) in a magnetic field, according to both numerical [9] and analytic [10] analysis. Deviations from the $SU(4)$ symmetry can be caused by variations of different-site interaction [3,5,11,12]. Another possibility for deviation was considered for an $SU(2) \otimes SU(2)$ Ising-type same-site anisotropy in Ref. [13,10], which may occur when the external field turns the spin oriented in parallel to the orbital angular momentum frozen in some direction by a crystalline field. A more generally-existing same-site interaction is $SU(3) \otimes U(1)$ one-ion L - S coupling (OILSC), the strength of which varies substantially in a range of $0 \sim 10^4 \text{cm}^{-1}$ ($1 \text{cm}^{-1} \sim 1 \text{K} \sim 10^{-4} \text{eV}$) for different elements [14]. As it is well understood now, L - S coupling can cause magnetic anisotropy [15], as far as a single ion is concerned. An interesting problem is to investigate the influence of such an OILSC on the collective properties, especially the QPT's of strongly-correlated spin-orbital systems in a magnetic field. It can be expected that the OILSC will bring about novel physics in the competition of spin and orbital degrees of freedom.

In the present paper we shall exactly solve a spin-orbital chain with such an OILSC via the Bethe ansatz (BA) approach. In the context of the different g factors of spin and orbit sectors, the OILSC leads to a rich variety of novel QPT's, with the analytic critical fields

explicitly obtained. Both spin ordering and orbital ordering exist in a gapped singlet phase, accompanied by a nonvanishing magnetization. The energy of the singlet is no longer invariant in the field due to the different g factors, which brings about five consecutive QPT's. It also contributes to unsymmetric changes of the total magnetization within a same phase in weak and in strong fields.

We consider an N -site chain with the Hamiltonian

$$\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_{L-S} + \mathcal{M}, \quad \mathcal{H}_0 = \sum_{i=1}^N P_{i,i+1},$$

$$\mathcal{H}_{L-S} = \lambda \sum_i \vec{l}_i \cdot \vec{s}_i, \quad \mathcal{M} = -g_s H \sum_i (g_s s_i^z + g_t l_i^z), \quad (1)$$

where $\vec{s}$ and $\vec{l}$ are spin-1/2 operators for spin and orbit and g_s and g_t are the corresponding Landé g factors, $P_{i,i+1} = (2\vec{s}_i \cdot \vec{s}_{i+1} + \frac{1}{2})(2\vec{l}_i \cdot \vec{l}_{i+1} + \frac{1}{2})$ exchanges the neighbor-site states. H is the magnetic field and periodic boundary conditions are applied throughout. $\mathcal{H}_{L-S}$ is the OILSC from the N sites and note that electrons have positive λ whereas holes have negative λ [15]. In the absence of the field, $\mathcal{H}_{L-S}$ breaks the system symmetry into $SU(3) \otimes U(1)$ when the bulk part $\mathcal{H}_0$ is $SU(4)$ invariant [6]. The basis consists of singlet and triplet of the $SU(2)$ Lie algebra $\{s^+ + l^+, s^- + l^-, s^z + l^z\}$, which is valid in the presence of the field only if $g_s = g_t$ as in the integrable spin ladder [16,17]. However, $\mathcal{H}_{L-S}$ and $\mathcal{M}$ do not commute due to the different g factors of spin and orbit, the conventional singlet and triplet are no longer their common eigenstates and fail to be the solution in the spin-orbital system. Nevertheless, we can diagonalize $\mathcal{H}_{L-S} + \mathcal{M}$ as a whole, which requires a new basis for all sites

$$\varphi_0 = \frac{\phi_1 - y^{-1}\phi_2}{\sqrt{1+y^{-2}}}, \quad \varphi_1 = \phi_3, \quad \varphi_2 = \frac{\phi_1 + y\phi_2}{\sqrt{1+y^2}}, \quad \varphi_3 = \phi_4,$$

where $\phi_1 = |\uparrow\downarrow\rangle$, $\phi_2 = |\downarrow\uparrow\rangle$, $\phi_3 = |\uparrow\uparrow\rangle$, $\phi_4 = |\downarrow\downarrow\rangle$ denote site state $|s^z l^z\rangle$ and $y^{\pm 1} = \pm g_- H / \lambda + \gamma(H/\lambda)$,

$\gamma(h) = (1 + g_-^2 h^2)^{1/2}$, $g_{\pm} = g_s \pm g_t$. The new realization of the SU(4) Lie algebra takes $S_{mn}\varphi_i = \delta_{ni}\varphi_m$,

$$\begin{aligned} S_{10} &= p(y)r(y)[yl^+(1/2 + s^z) - s^+(l^z + 1/2)], \\ S_{30} &= p(y)r(y)[ys^-(1/2 - l^z) - l^-(1/2 - s^z)], \\ S_{12} &= r(y)[l^+(1/2 + s^z) + ys^+(1/2 + l^z)], \\ S_{32} &= r(y)[s^-(1/2 - l^z) + yl^-(1/2 - s^z)], \quad S_{13} = s^+l^+, \\ S_{02} &= p(y)r^2(y)[y^2s^+l^- - s^-l^+ + y(s^z - l^z)], \end{aligned} \quad (2)$$

where $r(y) = 1/\sqrt{1+y^2}$, $p(y) = y/|y|$, other operators are their conjugates $S_{nm} = S_{mn}^\dagger$. For a certain choice of the basis order, the three Cartan operators I_k ($k = 1, 2, 3$) can be generated by commutations, e.g., $I_1 = [S_{23}, S_{32}]$, $I_2 = [S_{12}, S_{21}]$ and $I_3 = [S_{01}, S_{10}]$ for basis order $(\varphi_0, \varphi_1, \varphi_2, \varphi_3)^T$. The new basis $\{\varphi_0\}$ and $\{\varphi_1, \varphi_2, \varphi_3\}$ form the singlet and triplet of a new SU(2) Lie algebra $\{A^+, A^-, A^z\}$ where $A^z = s^z + l^z$ and

$$A^\pm = \sqrt{2}r(y)[(y-1)(s^\pm l^z - l^\pm s^z) + \frac{y+1}{2}(s^\pm + l^\pm)],$$

though the field breaks the symmetry into four U(1)'s.

We notice that a time-dependent single L - S coupled particle was solved by an SU(2) gauge transformation [18], but here we consider a many-body system with N correlated ions. Note that $P_{i,j}$ is still the permutation operator in the new basis, and $\mathcal{H}_{L-S} + \mathcal{M}$ commutes with the SU(4) bulk part $\mathcal{H}_0$. Based on the above new SU(4) realization we can exactly solve the model via the BA approach [8] with the BA equations and the eigenenergy

$$\begin{aligned} - \prod_{m=1}^{M^{(k)}} \Xi_1(\mu_{j,m}^{k,k}) &= \prod_{m=1}^{M^{(k+1)}} \Xi_{\frac{1}{2}}(\mu_{j,m}^{k,k+1}) \prod_{m=1}^{M^{(k-1)}} \Xi_{\frac{1}{2}}(\mu_{j,m}^{k,k-1}), \\ E &= - \sum_{j=1}^{M^{(1)}} 2\pi a_1(\mu_j^{(1)}) + \sum_{i=1}^4 E_i N_i, \end{aligned}$$

where $\Xi_x(\mu_{j,m}^{k,l}) = (\mu_j^{(k)} - \mu_m^{(l)} - xi)/(\mu_j^{(k)} - \mu_m^{(l)} + xi)$, $\mu_j^{(0)} = 0$, $M^{(0)} = N$, $M^{(4)} = 0$, and $1 \leq k \leq 3$; $a_n(\mu) = \frac{1}{2\pi} \frac{n}{\mu^2 + n^2/4}$, $E_0 = -\frac{\lambda}{4} - \frac{\lambda}{2}\gamma(H/\lambda)$, $E_1 = \frac{\lambda}{4} - \frac{1}{2}g_+H$, $E_3 = \frac{\lambda}{4} + \frac{1}{2}g_+H$ and $E_2 = -\frac{\lambda}{4} + \frac{\lambda}{2}\gamma(H/\lambda)$. N_i is the total site number in state φ_i . The basis order is chosen as $(\varphi_{P_1}\varphi_{P_2}\varphi_{P_3}\varphi_{P_4})^T$, where $P_i \in \{1, 2, 3, 4\}$ and φ_{P_1} is energetically the most favorable while φ_{P_4} is the least favorable. The relation $N_{P_i} = M^{(i-1)} - M^{(i)}$ gives another form of energy $E = \sum_{j=1}^{M^{(1)}} g^{(1)}(\mu_j) + g^{(2)}M^{(2)} + g^{(3)}M^{(3)}$. The field shifts the order from $(\varphi_0, \varphi_1, \varphi_2, \varphi_3)^T$ to $(\varphi_1, \varphi_0, \varphi_2, \varphi_3)^T$ for $\lambda > 0$, and $(\varphi_1, \varphi_2, \varphi_3, \varphi_0)^T$ to $(\varphi_1, \varphi_2, \varphi_0, \varphi_3)^T$ for $\lambda < 0$, at point $H_R = g_+ |\lambda| / (2g_s g_t)$. Following Ref. [19] and Ref. [20] one can obtain the ground state (GS) equations for three dressed energy $\epsilon^{(i)}$,

$$\epsilon^{(i)} = g^{(i)} - a_2 * \epsilon^{(i)-} + a_1 * (\epsilon^{(i-1)-} + \epsilon^{(i+1)-}), \quad (3)$$

where $\epsilon^{(0)} = \epsilon^{(4)} = 0$ and the symbol $*$ denotes the convolution. Fermi seas filled by negative $\epsilon^{(i)-}$ form the GS.

In the absence of the field, the GS is composed of all the four basis components when $\lambda_{c-} < \lambda < \lambda_{c+}$ where $\lambda_{c+} = 4$, $\lambda_{c-} = -2a_0/3$ and $a_0 = (\sqrt{3}\pi - 3\ln 3)/2$. But a strong positive OILSC $\lambda > \lambda_{c+}$ keeps the triplet gapful, the GS only consists of singlet and the system exhibits spin and orbital ordering, as shown later on. The first QPT will occur when the field brings down the triplet φ_1 and closes the gap $\Delta = g^{(1)}(0)$ at a critical field

$$H_{c0} = \frac{\lambda - 8 + \sqrt{\lambda^2 - \Delta_g^2 16(\lambda - 4)}}{g_+(1 - \Delta_g^2)} \quad (4)$$

where $\Delta_g = g_-/g_+$. Further increase of the field will exhaust all the singlet and so fully polarize the GS at

$$H_f = \frac{\lambda + 8 + \sqrt{\lambda^2 + \Delta_g^2 16(\lambda + 4)}}{g_+(1 - \Delta_g^2)}, \quad (5)$$

which is valid for all values of λ . For a weak OILSC $0 < \lambda < \lambda_{c+}$, the field brings out the components $\varphi_3, \varphi_2, \varphi_0$ in turn at three different critical points.

For $\lambda < 0$, the single-ion energy E_0 of the singlet φ_0 rises when the field becomes stronger. The triplet component φ_3 has lower energy in zero field but will be raised by the field in a quicker way than φ_0 . The energy competition of φ_3 and φ_0 results in new kinds of QPT's. In the four-component GS a weak negative OILSC does not expel the singlet far away from the triplet. In an increasing field φ_3 rises in energy more quickly and soon goes beyond φ_0 , so φ_3 gets out of the GS before φ_0 . There are three critical fields consecutively exhausting φ_3, φ_0 and φ_2 from the GS. However, a stronger negative OILSC leaves a longer distance between the triplet and singlet. Before φ_3 covers this distance to φ_0 , the singlet φ_0 has already risen completely out of the GS, which happens in the first critical point. But φ_3 soon draws very near to φ_0 such that φ_0 becomes close enough to the GS, in which φ_3 still lies, and is drawn back into the GS. This brings about the second QPT. Afterwards, the field overwhelms the influence of OILSC and pump out φ_3, φ_0 and φ_2 one by one, which gives other three QPT's. Therefore the system undergoes five consecutive QPT's (5-QPT) in this case, the variations of the components in the GS are: $1230 \rightarrow 123 \rightarrow 1230$ (and 1203) $\rightarrow 120 \rightarrow 12 \rightarrow 1$, where we denote the component φ_i by number i and each arrow indicates a QPT. Such unusual 5-QPT occur near λ_{c-} ($\lambda_{5c} < \lambda < \lambda_{c-}$, $\lambda_{5c} \doteq -0.712$, for $g_s = 2.0$, $g_t = 1.0$). When the OILSC is negatively stronger than J_{c-} , the singlet is gapful and does not exist in the GS before the field is applied. But the quicker rising of φ_3 will get the singlet involved in the GS, as what happens in the second QPT of the 5-QPT. In such case, four consecutive QPT's (4-QPT) will be observed: $123 \rightarrow 1230$ (and 1203) $\rightarrow 120 \rightarrow 12 \rightarrow 1$. A negatively stronger OILSC than

λ_Q will expel the singlet too far away from the GS composed of the triplet. The triplet component φ_3 itself will get out of the GS before it becomes adhesive and draw the singlet into the GS. We then have only the two QPT's.

We present a detailed phase diagram in Fig.1, the inset illustrates the 5- and 4-QPT. 5-QPT exist if C_-Q tends to rise up at C_- , with requirement on the g factors $g_t < (1 - \Delta_{g5})g_s/(1 + \Delta_{g5}) \doteq 0.516g_s$, where $\Delta_{g5} \doteq 3[(1+w)a_0]^{1/2}/(4\pi)$ and $w = 2/3$, satisfied for most typical spin-orbital systems. When the triplet φ_3 attracts the singlet φ_0 into the GS, some density of φ_3 themselves are attracted upwards a bit by φ_0 , which causes a quicker increase in magnetization. This attraction-and-counter-attraction is more sensitive near the point Q where φ_3 has small proportion in the GS. A 4-QPT magnetization near point Q is plotted in Fig.2B. The magnetization in the 5-QPT case has less apparent changes in the first two transition points due to heavier density of φ_3 .

Besides the exact singlet phase critical point (4) and full-polarized point (5), other critical points of the QPT can be obtained from Wiener-Hopf method [21]

$$\begin{aligned} H_c^{C+NQ} &\doteq H_+ + \tau k_+ G_{-1,1,-1}^2(\lambda, H_+), \\ H_c^{QD} &\doteq H_- + \tau k_- G_{-1,1,-1}^2(\lambda, H_-), \\ H_c^{C+M} &\doteq H_0 + \tau k_0 [w G_{\frac{1}{2}, -\frac{1}{2}, \frac{3}{2}}^2(\lambda, H_0) + G_{-\frac{1}{2}, \frac{1}{2}, \frac{3}{2}}^2(\lambda, H_0)], \\ H_c^{MQ} &\doteq H_0 + \tau k_0 [w G_{-1,1,0}^2(\lambda, H_0) + G_{-\frac{1}{2}, \frac{1}{2}, \frac{3}{2}}^2(\lambda, H_0)], \end{aligned}$$

where $k_0 = 3g_+^{-1}$, $\tau = 1/(4\pi^2)$, $H_0 = (a_0 - \lambda/2)g_+^{-1}$,

$$\begin{aligned} H_{\pm} &= \frac{\lambda_{\pm} - 3^{\pm 1} \sqrt{(\lambda_{\pm}^2 - 3^{1\pm 1} \lambda^2) \Delta_g^2 + \lambda^2}}{3^{(1\mp 1)/2} g_+ (1 - 9^{\pm 1} \Delta_g^2)}, \\ G_{l,m,n}(\lambda, H) &= l\lambda/2 + mg_+ H/2 + n \sqrt{\lambda^2 + g_-^2 H^2}/2, \\ k_{\pm} &= \frac{4}{3^{(1\mp 1)/2} g_+} \left(1 + \frac{3^{(1\pm 3)/2} \Delta_g^2 g_+ H_{\pm}}{\lambda_{\pm} - 3^{(1\mp 1)/2} g_+ H_{\pm}} \right)^{-1}, \end{aligned}$$

and $\lambda_{\pm} = 8 \ln 2 \pm \lambda$. $H_{\pm}$ and H_0 are the contributions of infinity Fermi points (FP), the other terms come from the revision of the finite FP. For $g_s = 2g_t$, H_+ can be simplified as $H_+ = [\lambda_+^2 - 9\lambda^2]/[2g_+\lambda_+]$. The above analytic results are compared with the numerical ones in Fig.1, which shows an excellent accuracy. For C_+NQD and C_+MQ , deviations only occur near C_+ due to small FQ invalid for Wiener-Hopf method, but an expansion based on small FP can be carried out and $H_c \cong (2c/\pi)g_+^{-1}(\lambda_{c+} - \lambda)^{3/2}$, $c = 1$ for C_+N and $c = (\ln 2)/3$ for C_+M . Setting $\lambda = 0$ in H_c^{MQ} , H_c^{C+NQ} and H_f will recover the very accurate expressions of the $SU(4)$ model obtained in Ref. [10].

A remarkable progress in experiments of spin-orbital systems is the observation of orbital ordering in some transition-metal compounds [2], which can be also found in the present model. The expectations of the spin

$\vec{s}$ and orbital angular momentum $\vec{l}$, the magnetization $M^z = g_s s^z + g_t l^z$ can be obtained for a singlet site

$$\langle \vec{s} \rangle_0 = -\langle \vec{l} \rangle_0 = g_- H [2\lambda\gamma(H/\lambda)]^{-1}, \quad \langle M^z \rangle_0 = g_- \langle \vec{s} \rangle_0.$$

Many transition-metal elements involve a strong OILSC and the GS of the system can be located in the singlet phase, in which all sites are occupied by the singlet. Consequently, the spins will be aligned ferromagnetically in the z direction while all the orbitals are ferromagnetically ordered in the opposite direction. Therefore, both orbital ordering and spin ordering can be observed. In addition, unlike the conventional singlet, the magnetization $\langle M^z \rangle_0$ is no longer zero in the field, as shown by the dash lines of curves e and f in Fig.2A. $\langle M^z \rangle_0$ increases almost in a linear way due to a small value of H/λ before the singlet phase is transited to another phase at H_{c0} . But the increasing rate $d\langle M^z \rangle_0/dH$ becomes smaller when the field gets stronger. This will lead to unsymmetric increasing of the magnetization in weak and in strong fields. The dotted regions of curves c , d and e in Fig.2A show a quicker climbing than the regions thereafter, which happens in a same phase with only two components φ_0 and φ_1 in the GS. Curve d only involves these two components from the beginning and provides a full view of this unsymmetry. Besides the slowdown of $d\langle M^z \rangle_0/dH$, another contribution to the unsymmetry comes from the variation in the energy-difference increasing rate. The energy difference $|E_1 - E_0|$ between the GS components φ_1 and φ_0 varies in a slower rate $d|E_1 - E_0|/dH$ when the field reaches higher.

We also present two typical magnetization for negative OILSC, shown by curves a and b in Fig.2A. The corresponding QPT's are: (a) 123 $\rightarrow$ 12 $\rightarrow$ 1; (b) 1230 (and 1203) $\rightarrow$ 120 $\rightarrow$ 12 $\rightarrow$ 1. One can notice the crossover of the two curves in phase 12 before full polarization, where only two components φ_1 and φ_2 are left in the GS. Curve b lies in four-component phase 1230 in the absence of the field, while a stronger OILSC expels the singlet off the GS and curve a starts from triplet phase 123. The negative λ makes φ_0 rising in energy level while φ_2 sinks when the field is applied. But the lowest state φ_1 sinks fast and so all the other components are moving out of the GS. Thus there is one more component φ_0 getting out of the GS for case b than case a . Therefore, the proportion of the most energetically favorable component φ_1 , with a positive magnetization $g_+/2$, increases more quickly in case b than in case a , which gives a quicker rise of magnetization in curve b before the first QPT. The component φ_3 in case b has smaller proportion in the beginning of a four-component phase 1230, while it has larger proportion in case a starting from a three-component phase 123. Consequently, φ_3 gets out of the GS earlier in case b and the first QPT occurs ahead of case a . But the rise of curve b soon slows down after its earlier first QPT, since the φ_0 has weaker negative magnetization than φ_3 in case a . After φ_3 gets out in case a , curve b soon rises

beyond curve a again as it has two components φ_2 and φ_0 moving out while only one component φ_2 in case a . As a result, case b has higher magnetization than case a when they both come to phase 12. But in this phase, φ_2 rises away from the lower state φ_1 always more quickly in case a with a stronger λ , as shown by $d|E_2 - E_1|/dH$ in Fig.2B. Therefore, φ_2 gets out faster in case a , and its lower magnetization rises in advance of case b and reaches the saturation point first. This results in the crossover.

Finally we stress only in the context of the different g factors of spin and orbit that the OILSC leads to these properties. If $g_s = g_t$ [17], the novel phenomena including the five consecutive QPT's, the spin and orbital ordering, nonzero magnetization of the singlet, the unsymmetric magnetization and the magnetization crossover, will all disappear.

We thank Huan-Qiang Zhou, You-Quan Li and Murray T. Batchelor for helpful discussions. ZJY thanks FAPERGS and FAPERJ for support. XWG thanks Australian Research Council. AF thanks FAPERGS and CNPq. IR thanks PRONEX and CNPq. BC thanks Zhejiang Natural Science Foundation RC02068.

[1] Y. Tokuro and N. Nagaosa, Science **288**, 462 (2000).




FIG. 1. Phase diagram of quantum phase transitions (QPT's) in magnetic field, $g_s = 2.0$, $g_t = 1.0$. The number i labels the state φ_i , e.g., the state components in phase 0123 are $\varphi_0\varphi_1\varphi_2\varphi_3$ in which the singlet φ_0 is energetically the most favorable whereas the triplet component φ_3 is the least favorable. The discrepancy between the analytic and numerical critical fields is not visible for most region of C_+NQD and C_+MQ . The boundary of singlet phase and the full-polarized point are exact. INSET: the five consecutive QPT's (dark dots) and four QPT's (dark boxes), their full-polarized points are relatively far away.

- [2] E. Saitoh *et al.*, Nature **410**, 180 (2001).
- [3] D.P. Arovas and A. Auerbach, Phys. Rev. B **52**, 10114 (1995).
- [4] A.V. Onufriev and J.B. Marston, Phys. Rev. B **59**, 12573 (1995).
- [5] S.K. Pati and R.R.P. Singh, Phys. Rev. Lett. **81**, 5406(1998).
- [6] Y.Q. Li, M. Ma, D.N. Shi and F.C. Zhang, Phys. Rev. Lett. **81**, 3527 (1998).
- [7] Y.Q. Li, M. Ma, D.N. Shi and F.C. Zhang, Phys. Rev. B **60**, 12781 (1999).
- [8] B. Sutherland, Phys. Rev. B **12**, 3795 (1975).
- [9] S.J. Gu and Y.Q. Li, Phys. Rev. B **66**, 092404 (2002).
- [10] Z.J. Ying, A. Foerster, X.W. Guan, B. Chen and I. Roditi, cond-mat/0308443.
- [11] F. Mila, B. Frischmuth, A. Deppeler and M. Troyer, Phys. Rev. Lett. **82**, 3697 (1999).
- [12] P. Azaria, A.O. Gogolin, P. Lecheminant and A.A. Nersisyan, Phys. Rev. Lett. **83**, 624 (1999).
- [13] S.J. Gu, Y.Q. Li and H.Q. Zhou, cond-mat/0308432.
- [14] R.D. Cowan, *The theory of atomic structure and spectra* (University of California Press, Berkeley, 1981).
- [15] C.P. Slichter, *Principles of magnetic resonance* (Springer-Verlag, New York, 1990).
- [16] Y. Wang, Phys. Rev. B **60**, 9236 (1999).
- [17] M.T. Batchelor, X.W. Guan, A. Foerster and H.Q. Zhou, New J. Phys. **5**, 107 (2003).
- [18] S.J. Wang and L.X. Cen, Phys. Rev. A **58**, 3328 (1998).
- [19] M. Takahashi, Prog. Theor. Phys. **46**, 401 (1971).
- [20] C.N. Yang and C.P. Yang, J. Math. Phys. **10**, 1115(1969).
- [21] M.G. Krein Usp. Mat. Nauk **13**, 3 (1958).




FIG. 2. (A) Typical magnetization behaviors in different QPT's, $g_s = 2.0$, $g_t = 1.0$. The dashed lines are gapped singlet phase with a nonzero magnetization M^z , both spin and orbital ordering can be observed in this phase. Curve d provides a full view of unsymmetric growth of M^z , the dotted-line region (phase 01) in weak field climbs faster than the solid-line region (phase 10) in strong field. Also note the crossover between curve a and b before full polarization. (B) Magnetization of four consecutive QPT's, $\lambda = -0.76$, $H_{c4} = 3.65$, the numbers denote the phases as in Fig.1. (C) Comparison on increasing speed of energy difference $E_2 - E_1$ in the field for $\lambda = -1.5$ and $\lambda = -0.5$, the former always has a larger speed.